

Pattern of Sensorineural Hearing Loss in Patients Attending ENT OPD

Manish Sharma¹, Padam Singh², Megha Kapoor³, Mohit Goel⁴

1- Senior Resident, Department of ENT and Head & Neck Surgery, Government Medical College, Jammu. 2- Associate Professor, Department of ENT and Head & Neck Surgery, Government Medical College, Jammu. 3- Post Graduate in Department of Physiology, Santosh Medical College, Ghaziabad, U.P. 4- Post Graduate in Department of ENT and Head & Neck Surgery, Government Medical College, Jammu.

Correspondence to:
Dr. Manish Sharma, Senior Resident, Department of ENT and Head & Neck Surgery, Government Medical College, Jammu.
Contact Us: www.ijohmr.com

ABSTRACT

Introduction: Hearing skills is one of the most important and pleasurable gifts bestowed by almighty which affects development of an individual. Sensorineural hearing loss (SNHL) is one of the types of hearing loss in which the root cause lies in the inner ear, vestibulocochlear nerves or central processory centres of the brain. It can be present at birth (congenital) or can develop later in life (delayed). **Materials and methods:** This study was conducted over a period of one year to find out the pattern of SNHL in the patients attending ENT OPD. A total of 820 cases with pure SNHL were analyzed. After detailed case history and ENT examination of the selected patients, audiological evaluation was performed in audiology room. Brain Evoked Response Audiometry (BERA) was performed in patients less than 5 yrs and Short Increment Sensitivity Index (SISI), Tone decay, Alternate Binaural Loudness Balance test (ABLB) was done in patients with pure SNHL in patients of age more than 5 yrs confirmed with Pure Tone Audiometry (PTA). **Results:** A total of 43003 patients visited ENT OPD and out of them 820 patients were diagnosed with pure sensorineural hearing loss (SNHL) with a incidence of 19.06 per thousand population and sex ratio of 1.35:1 (male: female). Maximum patients were in the age group of 55-75 (64.27%). Gradual hearing loss was found in 778 (95%) patients and 42 (5%) patients presented with sudden sensorineural hearing loss (SSNHL) with bilateral involvement in 675 (82.3%) and unilateral in 145 (17.6%) patients. **Conclusion:** Presbycusis was found to be the most common cause of SNHL in patients of age more than 15 yrs and in patients less than 15 yrs, no apparent cause was found in most of the patients.

KEYWORDS: ABLB, BERA, Presbycusis, PTA, Sensorineural hearing loss, SISI

INTRODUCTION

Hearing skills is one of the most important and pleasurable gifts bestowed by almighty which affects the development of an individual. Sensorineural hearing loss (SNHL) is a type of hearing loss in which the root cause lies in the inner ear, vestibulocochlear nerves or central processory centres of the brain. It can be present at birth (congenital) or can develop later in life (delayed). It can be genetic or result from acquired factor such as disease or injury. The incidence of bilateral SNHL ranges from 1.4 to 3 per thousand live births in various studies worldwide.¹⁻⁴ Of the SNHL described in children, 50% is thought to be genetic, 25% acquired and 25% of unknown etiology.³ In children, SNHL may accompany syndromes such as:- Waardenburg's syndrome, Albinism, Pendred syndrome, Down's syndrome. Acquired causes of SNHL in children include Maternal use of ototoxic drugs, Viral infections that affect the fetus like rubella, Metabolic disorders, Rh incompatibility, Irradiation in the first trimester and Idiopathic. In adults, the most important risk factor for SNHL is age.⁵ Other factors are Noise induced, Ototoxicity, Idiopathic sudden SNHL, Meniers disease, Trauma, Syphilis, Lyme disease etc..

MATERIALS AND METHODS

This study was conducted in the Department of ENT &

Head and Neck Surgery, SMGS Hospital, Government Medical College, Jammu, during the year 2009 to 2010 to study the role of different etiological factors associated with SNHL and to study the age and sex wise distribution of cases presenting with SNHL. A total of 820 cases with pure SNHL were analyzed for this study from patients attending ENT OPD. Case selection was based on following criteria:

- Cases without any previous history of ear discharge and normal tympanic membrane on Otoscopy.
- Cases with pure SNHL (conductive and mixed hearing loss patients excluded), whether congenital or acquired as detected by pure tone audiometry (In patients with age more than 5 yrs) and BERA (In patients with age less than 5 yrs).

After proper consent, detailed case history, and ENT examination, an audiological evaluation was performed in audiology room. BERA was performed in patients less than 5 yrs and SISI, Tone decay, ABLB was done in patients with pure SNHL in patients of age more than 5 yrs confirmed with PTA. Other investigations include examination of blood, urine and MRI of internal acoustic meatus whenever essential.

How to cite this article:

Sharma M, Singh P, Kapoor M, Goel M. Pattern of Sensorineural Hearing Loss in Patients Attending ENT OPD. *Int J Oral Health Med Res* 2015;2(1):5-8.

RESULTS

(A) The incidence of SNHL -: The data collected from hospital records, between 2009 to 2010, revealed that SNHL was 19.06 per thousand population of all the patients who attended ENT OPD. Out of total 820, maximum cases were seen in the age group of 55-75 yrs (64.27%), and least number of patients were seen in the age group of 45-55yrs (2.68%). 52 patients (6.34%) were below the age of 15 yrs. A total of 472 (58%) cases were males and 348 (42%) cases were females with male to female sex ratio of 1.35:1. The age-sex distribution is shown in table 1. Out of 820 patients, those having bilateral hearing loss were 675 (82.3%) and unilateral hearing loss were seen in 145 (17.6%) patients. Amongst unilateral cases, right ear involvement was in 48.2% cases and left ear involvement was in 51.8%, shown in table 2.

Age(yrs)	No. of patients	Percentage (%)	Males	Females	Sex ratio
<15	52	6.34	34	18	1.88:1
15-45	87	10.61	43	44	0.97:1
46-55	22	2.68	17	5	3.4:1
56-75	527	64.27	303	224	1.35:1
>75	132	16.1	75	57	1.31:1
Total	820	100	472	348	1.35:1

TABLE 1: AGE SEX DISTRIBUTION (n=820)

Laterality	No. of patients	Right ear	Left ear
Bilateral	675 (82.3%)		
Unilateral	145 (17.6%)	70 (48.2%)	75 (51.8%)
Total	820 (100%)		

TABLE 2: MONOAUROAL OR BINAURAL (n=820)

(B)The severity of SNHL -: In our study of 820 patients, 260 patients (36.8%) were having severe hearing loss and only 37 patients have mild hearing loss as shown in figure 1.




FIGURE 1: PERCENTAGE OF HEARING IMPAIRMENT (n=820)

(C) Sudden or gradual -: In our study of 820 patients, 42 (5%) presented with sudden hearing loss and 778 (95%) presented with a gradual hearing loss.

(D) Analysis of cases of SNHL in patients with age > 15 yrs (n=768)

Etiology- Probable etiology of SNHL was found in 726 cases (94.5%), and 42 cases (5.49%) were idiopathic. Out

of these 726 cases, 660(85.9%) cases were of Presbycusis, followed by Noise induced in 14 cases (1.82%) and systemic diseases in 29 cases (3.77%). Other causes were Trauma, Otosclerosis, Ototoxic drugs, Vestibular schwannoma and Meniers disease.

Age distribution- Table 3 shows the distribution of etiology in different age groups. This indicates that Presbycusis starts after the age of 45 yrs. Out of 660 cases of Presbycusis, 420 (64%) were in the age group of 55-75 yrs. All the cases of noise trauma were between the age group of 15-45 yrs.

Laterality- In our study, 627 patients (82%) were having both ears involved and in 141 cases (18%), only one ear was involved.

Etiological factors	No. of patients	% age	U/L	B/L	15-45 yrs	45-55 yrs	55-75 yrs	>75 yrs
Presbycusis	660	85.93	111	549	0	90(14%)	420(64%)	200(30%)
Noise	14	1.82	1	13	14(100%)	0	0	0
Trauma	6	0.78	6	0	2(33%)	0	3(50%)	1(17%)
Otosclerosis	2	0.26	2	0	2(100%)	0	0	0
Ototoxic drugs	4	0.52	3	1	4(100%)	0	0	0
Vestibular schwannoma	1	0.13	1	0	0	0	1(100%)	0
Meniers disease	10	1.3	5	5	0	8(80%)	2(20%)	0
Systemic diseases	29	3.77	2	27	0	9(31%)	12(41%)	8(28%)
Idiopathic	42	5.49	10	32	12(29%)	12(29%)	10(24%)	8(18%)
Total	768	100	141(18%)	627(82%)				

TABLE 3: ETIOLOGICAL FACTORS OF SNHL IN PATIENTS WITH AGE > 15 YRS (n=768):

Apparent etiological factors	No. of patients	%age	U/L	B/L
Prenatal	5	9.7		5
Intranatal	4	7.6		4
Postnatal	12	23.2	2	10
Congenital syndromic	14	26.9		14
No apparent cause	17	32.6	2	15
Total	52	100	4	48

TABLE 4: APPARENT ETIOLOGICAL FACTORS IN CHILDREN < 15 YRS (n=52)

(D) Analysis of cases of SNHL in patients with age < 15 yrs (n=52)

Etiology- No apparent cause could be established in 32.6% cases. Deafness due to etiological factors in the postnatal period was seen in 23.2% cases, prenatal period in 9.7% cases and intranatal period in 7.6% cases. SNHL associated with congenital syndromes was proven to be in 26.9% of cases and out of them Down's syndrome was the most common syndrome as shown in table 4.

Age distribution- Amongst these 52 cases, 40 children presented with congenital SNHL and 12 patients presented later on.

Laterality- Out of 52 patients, 48 patients presented with bilateral hearing loss and only 4 patients with unilateral.

DISCUSSION

Incidence of SNHL- In our study the incidence of SNHL is 19.06 per 1000 population which is slightly higher as compared to Mathers C. et al. (2001)⁶ in which they studied global burden of hearing loss in 2000 and reported an age standardized incidence of 3.63 per 1000 males and 3.60 in female patients with an average of 7.1 patients of SNHL per 1000 population. Our higher incidence may be because of difference between the total populations being considered; we have taken only patients attending ENT OPD, and they have taken patients globally.

Age distribution - According to our study, out of 820 patients, the majority were in age group of 55-75 yrs (64%). Number of cases below 15 yrs were 52 (6%), and 132 cases (16%) were below 75 yrs of age. Our results are almost consistent with J.A.E.Eziyi.et al. (2009)⁷ who found in their study of 530 SNHL patients attending Nigerian Audiology clinic that maximum load of SNHL patients were between age group of 55-75 yrs (52.1%) and few patients (18.1%) were seen above 75 yrs of age.

Monaural or binaural- According to our study, out of 820 SNHL patients, 18% presented with unilateral hearing loss and 82% presented with bilateral hearing loss, which is consistent with that of Sculerati N (2000)⁸ who in a cohort study of 168 patients, found that SNHL was bilateral in 82% and unilateral in 18% cases. Out of 145 patients of unilateral SNHL, in our study, 48.2% patients presented with right ear involvement and 51.8% with left ear involvement which was consistent with the findings of PE Brookhouser et al. (1991)⁹ who in their study of 324 children and adolescents with unilateral SNHL at the Boys Town Research Hospital, found that, left ear was affected in 52% and right ear in 48%.

Sex wise distribution- In our study, out of 820 patients, 472 were males and 348 females with a sex ratio of 1.35:1. Yanagita N.et al. (1994)¹⁰ in their nationwide survey of SNHL in 1987 in Japan found the sex ratio of 1.1:1. The higher sex ratio in our study is because of the fact that the overall incidence of female patients in ENT OPD is less.

Hearing impairment - Most of the patients in this study had moderately severe (29.3%) and severe SNHL (36.8%) with profound hearing loss in 14% of cases whereas Magnus S.K. Johansson and Stig D. Arlinger (2003)¹¹ in a study of patients in Sweden of age group of 20-30 yrs, maximum patients were of mild hearing loss.

Onset of hearing loss - In our study of 820 patients, 42 (5%) presented with sudden hearing loss and 95% with gradual hearing loss in contrast to Hughes GB et al. (1996)¹² who, according to their study labeled that amongst all cases of SNHL in United States, approximately 1% of cases are of sudden SNHL. Maximum patients of sudden SNHL in our study

presented with unilateral hearing loss (97.7%) and only 1 patient (2.3%) presented with bilateral sudden SNHL. Our values are consistent with the study of Bruce L. Fetterman MD et al. (1996)¹³ in which they concluded that out of 823 patients of sudden SNHL, 14 (1.7%) had sudden bilateral SNHL and with the study of Yanagita N and Murahashi K (1987)¹⁴ who reported 10 patients with simultaneous bilateral SNHL out of 997(1%).

Etiological factors in patients of age > 15 yrs - Presbycusis was found to be the most common cause of SNHL in patients with age > 15 yrs. Similar findings are reported in other studies also. Mathers C. et al. (2001)⁶ and K H Yiap et al. (1984)¹⁵ reported that the leading causes of adult onset hearing loss are presbycusis followed by the noise induced hearing loss. Maximum number of patients with presbycusis were seen in the patients of age more than 55 yrs which is consistent with the results of Mosicki JK et al. (1985)¹⁶ who stated that prevalence of hearing impairment (presbycusis) is high as 83% in people between the age of 57 and 89 and Schoenborn CA and Marano M (1987)¹⁷ who also reported Presbycusis more in patients of age more than 65 yrs in United States. Systemic diseases like Diabetes (commonest), hypertension and thyroid dysfunction were found to be the cause of SNHL in 29 patients out of 768 adult patients (3.7%). Bainbridge K et al. (2009)¹⁸ reported the risk for SNHL in persons with diabetes. Noise induced hearing loss (NIHL) was noticed in 14 cases (2%), and all the 14 cases were between the age group of 15-45 yrs. In their study of Bangladesh textile factory workers, Md Yusuf Haider et al. (2008)¹⁹ resulted that incidence of NIHL was 20.59% in workers aged less than 35 yrs and it was 41.38% in age group above 35 yrs. Susan L. Phillips et al. (2010)²⁰ in their study amongst student musicians of age 18-25 yrs (n=329), concluded that the overall prevalence of NIHL was 45%. These studies show that there is age predisposition of NIHL and is more prevalent in the early age group of less than 45 yrs. These findings were consistent with our findings since NIHL cases in our study were in age group of 15-45 yrs. The higher prevalence of NIHL in the age group of 15-45 yrs can be explained on the fact that this is the age of teenagers hearing loud music and factory employees working in loud, noisy atmospheres. Other etiological factors for SNHL were Trauma, Otosclerosis, Ototoxicity, Vestibular schwannoma, Meniers disease and Idiopathic.

Etiological factors in patients of age < 15 yrs - In our study idiopathic SNHL is the commonest known cause of deafness in children less than 15 (33%). Other apparent causes of SNHL in children included postnatal (meningitis, neonatal jaundice, trauma, ototoxicity) in 23.2% cases, intranatal (birth asphyxia) in 7.6% cases, prenatal (maternal infections, ototoxic drugs taken by mother, radiation exposure of mother during 1st trimester) in 9.7% cases and syndromic SNHL in 30% of cases. Most of them were having bilateral SNHL and only 4 patients had a unilateral hearing loss. Dunmade AD et al. (2007)²¹ in their study found that in about a third (34.8%) of patients, causes were unknown, probably

congenital and the main acquired causes were febrile illness (18.3%), measles (13.9%), meningitis (8.7%), mumps (6.9%), or severe birth asphyxia (4.3%). Morzaria et al. (2004)²² found in their study of children less than 18 yrs of age, that, most common cause of bilateral SNHL are unknown (37.7%), prenatal (12%), perinatal (9.6%), postnatal (8.2%), genetic syndromic (3.2%) and genetic nonsyndromic. The higher number of cases without any apparent cause might be because of the higher number of cases of congenital deafness in our population which goes undiagnosed. This is supported by the fact that 27 children out of 52 in our study were the products of consanguineous marriages known to be a risk factor for congenital deafness as related by Reddy M.V.V et al. (2006)²³. Our study shows that syndromic patients were 30% and nonsyndromic 70% whereas Monisha Mukherjee et al. (2003)²⁴ in their study found, syndromic and nonsyndromic patients to be 40% and 60% respectively.

CONCLUSION

- a) A total of 820 patients of pure SNHL were diagnosed in a period of one year in the department of Otorhinolaryngology and head and neck surgery, SMGS hospital out of 43004 patients who attended ENT OPD with an incidence of 19.06 per 1000 population per year with male: female sex ratio of 1.35:1. The highest numbers of patients were seen in the age group of 55-75 yrs.
- b) Maximum cases were having bilateral ear involvement (82.3%), and unilateral ear involvement was less (17.6%).
- c) The gradual hearing loss was seen in 95% cases and only 5% presented with sudden sensorineural hearing loss (SSNHL).
- d) Most cases presented with severe (36.8%) and moderately severe (29.3%) hearing loss. e) Presbycusis was found to be the most common cause of SNHL in patients more than 15 yrs with maximum falling in the age group of 55-75 yrs followed by the idiopathic group.
- f) All the patients of NIHL were between the age group of 15-45 yrs.
- e) In maximum cases of SNHL patients aged less than 15 yrs, no apparent cause was found, followed by syndromic children and amongst those syndromic children, Downs syndrome was the most commonly found syndrome.

REFERENCES

1. White KR et al. Screening all newborns of hearing loss using transient evoked Otoacoustic emissions. *Internat J Pediatr Otolaryngol* 1994;29(3):203-217.
2. Mason JA et al. Universal infant hearing by automated auditory brainstem response measurement. *Pediatrics* 1998;101(2):221-228.
3. Markes HA et al. Congenital abnormalities of inner ear, disease of the ear, Bath, England: Arnold and Oxford University Press 1998;208-296.
4. Hone SW et al. Medical evaluation of pediatric hearing loss, laboratory radiographic and genetic testing. *Otolaryngol Clin Nor Amer* 2002;35(4):751-764.
5. George A. Gates et al. Incidence of hearing decline in elderly. *Acta Otolaryngol (Stockh)* 1991;111:240-248.
6. Mathers C et al. Global burden of hearing loss in the year 2000. *Global burden of disease 2000; World health report 2001*(2).
7. J.A.E. Eziyi et al. Audiological pattern of hearing loss at Obafemi Awolowo University Teaching Hospital Complex-Ile-Ife, Nigeria. *The Inter J Otolaryngol* 2009;8(2).
8. Sculerati N. Analysis of a cohort of children with sensory hearing loss using the scale systemic nomenclature. *Laryngoscope* 2000;110(5.1):787-98.
9. P E Brookhouser et al. Unilateral hearing loss in children. *Laryngoscope* 1991;101(12.1):1264- 1272.
10. Yangita N et al. Estimated annual number of patients treated for sensorineural hearing loss in Japan. *Acta Otolaryngol (Stockh)* 1994;Suppl 514:9-13.
11. Magnus S.K. Johansson and Stig D. Arlinger. Prevalence of hearing impairment in a population in Sweden. *Int J Aud* 2003;42:18-28.
12. Hughes GB et al. Sudden Sensorineural hearing loss. *Otolaryngol Clin North Am* 1996 Jun;29(3):393-405.
13. Bruce L. Fetterman MD et al. Sudden Bilateral Sensorineural hearing loss. *Laryngoscope* 1996;106(11):1347-1350.
14. Yangita N et al and Murahashi K. Bilateral simultaneous sudden deafness. *Arch Otorhinolaryngol* 1987;244:7-10.
15. K H Yiap et al. Hearing sensitivity of Singaporeans in advance age. *Singapore Medical Journal* 1984;25(6):420-423.
16. Mosicki JK et al. Hearing loss in the elderly: an epidemiological study of the Framingham heart study cohort. *Ear Hear* 1985;6:184-190.
17. Schoenborn CA and Marano M. Current estimates from the national health interview survey: United States 1987, *Vital and Health Statistics*; 10(6).
18. Bainbridge K et al. Hearing impairment an under-recognized complication of diabetes, *Diabetes Voice* March 2009;54(1).
19. Md Yusuf Haider et al. Noise induced hearing loss amongst the textile industry workers. *Bangladesh J Otorhinolaryngol* 2008;14(2):39-45.
20. Susan L. Phillips et al. Prevalence of noise induced hearing loss in student musicians. *Int J Audio* 2010 April;49(4):309-316.
21. Dunmade AD et al. Profound Sensorineural hearing loss in Nigerian children. Any shift in etiology? *J of Deaf Studies and Deaf Education* 12: (1) winter 2007.
22. Morzaria et al. Systemic review of the etiology of bilateral sensorineural hearing loss in children. *Int J Ped Otorhinolaryngol* 2004;68(9):1193-8.
23. Reddy M.V.V et al. Role of consanguinity in congenital neurosensory deafness; *Kamla-Raj* 2006 *Int J Hum Genet*, 2006;6(4):357-358.
24. Monisha Mukherjee et al. Connexin 26 and autosomal recessive nonsyndromic hearing loss. *Indian J Hum Genet* 2003;9(2):40-50.

Source of Support: Nil
Conflict of Interest: Nil